arXiv:1304.0607v1 [cond-mat.stat-mech] 2 Apr 2013

The three site interacting spin chain in staggered field: Fidelity vs Loschmidt echo

Uma Divakaran[1]

[1]Department of Physics, Indian Institute of Technology Kanpur, Kanpur 208 016, India

We study the fidelity and Loschmidt echo of a three site interacting XX chain in presence of staggered field which consists of special types of quantum phase transition points due to change in the topology of the Fermi surface, apart from normal phase transition points from gapped to gapless phases. We focus on the effect of such types of phase transition points on the fidelity, which is the static part of the problem, and Loschmidt echo of this spin chain. We find that on one hand, fidelity is able to detect only the boundaries of the gapped to gapless phase transition and not the phase transition from two Fermi points to four Fermi points which lie within this gapless region. On the other hand, Loschmidt echo is just able to detect one special point in the entire phase diagram. We present the arguments for such a behavior.

PACS numbers:

I. INTRODUCTION

The desire of building a quantum computer to solve quantum problems efficiently has lead to the development of quantum information theory and other related fields. Fidelity, which is basically an overlap of the ground state wave function at two slightly different values of the parameter, is an important component of the quantum information theory[1–3]. It is important because it can detect a quantum critical point without knowing anything about the order parameter of the system, which otherwise is the conventional way of looking at the quantum phase transition (QPT)[4,5]. Fidelity has been successful in detecting various types of quantum phase transition points, for example, ordinary critical points separating two gapped phases through a gapless point[1,2] or topological QPTs[6] or QPT in Bose Hubbard model[7], but its usefulness in a general scenario is not very clear. We present a model where the fidelity detects the onset of quantum critical region but not the two different phases within this quantum critical region.

One of the other important quantities in quantum information theory is Loschmidt echo (LE)[8–10]. LE is the measure of the overlap of the same initial state, the ground state of the initial Hamiltonian $H(\lambda)$, evolving under the influence of the two Hamiltonians, $H(\lambda)$ and $H(\lambda+\delta)$. This overlap shows a dip at the quantum critical point, thus enabling us to detect a quantum critical point. In the language of quantum information theory, LE can be used to measure the quantum to classical transition of a qubit coupled to an environment making it an important component of quantum information theory. LE was actually introduced in connection to the quantum to classical transition in quantum chaos[11–15] and now extended to various other systems like Ising model[8], Bose-Einstein condensate model[16] and Dicke model[17]. It has also been studied experimentally using NMR[18–20]. Once again, LE has been used to detect the quantum critical points but its generality is yet to be explored.

In this paper, we point out the inability of fidelity or LE to detect certain special class of critical points by taking the example of staggered transverse field in a three spin interacting spin-1/2 XX chain. This study is an important addition to the literature as no one has yet looked at the fidelity and LE of a Hamiltonian which also undergoes a very unique type of quantum phase transition, namely, from two Fermi points to four Fermi points. It is not apriori obvious the effect of such points on the fidelity and LE of the model. It is to be noted that many interesting experimental observations, like magnetic properties of solid ^{3}He, were explained by the inclusion of multispin interaction in the Hamiltonian[21].

The outline of the paper is as follows: The model is described briefly in Section II. Section III discusses the statics, i.e., the ground state fidelity of the model and the dynamics of ground state Loschmidt echo is studied in Section IV. We present our conclusions in Section V.

II. MODEL

In this section, we quickly go through the ground state phase diagram of the three spin interaction Hamiltonian in presence of a staggered field h_s given by the Hamiltonian,[22]

$$\begin{aligned} H_{TS} &= -\sum_j J(\sigma_j^x\sigma_{j+1}^x + \sigma_j^y\sigma_{j+1}^y) - J_3(\sigma_j^x\sigma_{j+2}^x + \sigma_j^y\sigma_{j+2}^y)\sigma_{j+1}^z \\ &- \sum_j (-1)^j h_s \sigma_j^z \end{aligned} \qquad (1)$$

Performing standard Jordan Wigner Fermionization[23,24], we get

$$\begin{aligned} H_{TS} &= -\frac{J}{2}\sum_j (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \frac{J_3}{4}\sum_j (c_j^\dagger c_{j+2} + c_{j+2}^\dagger c_j) \\ &- (-1)^j h_s (c_j^\dagger c_j - \frac{1}{2}) \end{aligned} \qquad (2)$$

To diagonalize the above Hamiltonian, it is useful to introduce two types of spinless fermions on the odd and even sub-lattices as shown below:

$$c_{2n-1} = a_{n-1/2} \text{ and } c_{2n} = b_n.$$

Substituting this in Eq. 2 and Fourier transforming it gives,

$$\begin{aligned} H_{TS} &= \sum_k H_k \\ &= \sum_k [\epsilon_a(k) a_k^\dagger a_k + \epsilon_b(k) b_k^\dagger b_k + \epsilon_{ab}(k)(a_k^\dagger b_k + b_k^\dagger a_k)] \end{aligned} \quad (3)$$

where

$$\begin{aligned} \epsilon_a(k) &= \frac{J_3}{2}\cos(k) + h_s \\ \epsilon_b(k) &= \frac{J_3}{2}\cos(k) - h_s \\ \epsilon_{ab}(k) &= -J\cos(k/2) \end{aligned} \quad (4)$$

It is easier to understand the fidelity and LE calculations if we rewrite the Hamiltonian H_k in the four basis described by $|0,0\rangle$, $|a_k,0\rangle$, $|0,b_k\rangle$, $|a_k,b_k\rangle$ where the first index represents the presence or absence of a_k- particle and the second index corresponds to that of b_k-particle. In these basis, H_k is given by

$$H_k = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \epsilon_a(k) & \epsilon_{ab}(k) & 0 \\ 0 & \epsilon_{ab}(k) & \epsilon_b(k) & 0 \\ 0 & 0 & 0 & \epsilon_a(k)+\epsilon_b(k) \end{bmatrix}. \quad (5)$$

It is clear from the above representation that to diagonalize the above Hamiltonian, only the basis $|a_k,0\rangle$ and $|0,b_k\rangle$ needs to be rotated since the Hamiltonian in the other two basis is already diagonal, i.e., there is no mixing along these two directions. Let us denote the two new directions for the two new quasi-particles α_k and β_k as $|\alpha_k,0\rangle$ and $|0,\beta_k\rangle$ which diagonalize the total Hamiltonian giving four eigen energies 0, $E_-(k)$, $E_+(k)$ and $E_-(k)+E_+(k)$ corresponding to the four directions $|0,0\rangle$, $|\alpha_k,0\rangle$, $|0,\beta_k\rangle$ and $|\alpha_k,\beta_k\rangle$, respectively. These two new directions are given by

$$\begin{aligned} |\alpha_k,0\rangle &= \cos\frac{\theta_k}{2}|a_k,0\rangle - \sin\frac{\theta_k}{2}|0,b_k\rangle \\ |0,\beta_k\rangle &= \sin\frac{\theta_k}{2}|a_k,0\rangle + \cos\frac{\theta_k}{2}|0,b_k\rangle \end{aligned} \quad (6)$$

where

$$\tan\theta_k(h_s) = \frac{-\cos k/2}{h_s} \quad (7)$$

with J set equal to unity through out the paper, and their energies are

$$E_\pm(k) = \frac{J_3}{2}\cos k \pm \sqrt{h_s^2 + \cos^2(k/2)}. \quad (8)$$

Hence, the Hamiltonian is now given by

$$H_{TS} = \sum_k E_-(k)\alpha_k^\dagger \alpha_k + E_+(k)\beta_k^\dagger \beta_k.$$

The ground state of this Hamiltonian corresponds to all the modes with negative energies filled and hence depending upon the values of the Hamiltonian parameter, the system has various phases. We discuss briefly the various phases below:



FIG. 1: The phase diagram of the three spin model in presence of staggered field. The solid lines correspond to $h_s = \pm J_3/2$ critical line and the dotted lines correspond to $h_s = \pm\sqrt{\frac{J_3^2-4}{4}}$.

- When $h_s > J_3/2$: $E_+(k) > 0$ and $E_-(k) < 0$ for all k, and hence the ground state for each mode is $|\alpha_k,0\rangle$ with total ground state energy $E_g = \sum_k E_-(k)$. This is the Antiferromagnetic (AF) phase.

- When $\sqrt{J_3^2/4-1} < |h| < J_3/2$: Some of the k modes from the $E_+(k)$ branch become negative whereas $E_-(k)$ is negative for all the modes. This phase has two Fermi points arising due to zeros of the $E_+(k)$ branch. In this region, the ground state for a given mode can be $|\alpha_k,0\rangle$ or $|\alpha_k,\beta_k\rangle$ depending upon whether $E_+(k)$ branch is empty or filled, respectively. The total ground state energy is given by

  $$\sum_k E_-(k)\Theta(-E_-(k)) + E_+(k)\Theta(-E_+(k)),$$

  where Θ is the Heaviside function. We call this phase as spin liquid I (SLI) phase.

- When $0 < |h| < \sqrt{J_3^2/4-1}$: In this limit, $E_-(k)$ also crosses zero for some modes and hence there are three possible ground states for a given mode k depending upon the signs of the energies $E_\pm(k)$ given by $|\alpha_k,0\rangle$, $|0,0\rangle$ and $|\alpha_k,\beta_k\rangle$. Once again the ground state energy is the sum over all the modes with negative energies for each branch as written above. This phase is called spin liquid II (SLII) phase.

Following the above arguments, the phase diagram of the model is shown in Fig. 1. We now try to capture these phase transitions using the fidelity approach, especially the phase transition from SLI to SLII and later discuss them in the light of Loschmidt echo.

III. FIDELITY

As mentioned in the introduction, the ground state fidelity is defined as the overlap between the two ground state wave functions at slightly different parameter values and is given by

$$F = \langle \Psi_G(h_s)|\Psi_G(h_s+\delta)\rangle = \prod_k \langle \Phi_k(h_s)|\Phi_k(h_s+\delta)\rangle = \prod_k F_k \tag{9}$$

where $|\Psi_G(h_s)\rangle$ is the total ground state for the given parameter value and $|\Phi_k(h_s)\rangle$ is the ground state for the $k-th$ mode. Thus, while evaluating F_k, one has to carefully identify the ground state for the $k-th$ mode depending upon the sign of $E_\pm(k)$. The various possibilities are:

- $F_k = \langle \alpha_k, 0|\alpha_k, 0\rangle_\delta = \cos(\frac{\theta_k(h_s)-\theta_k(h_s+\delta)}{2})$
- $F_k = \langle \alpha_k, \beta_k|\alpha_k, \beta_k\rangle_\delta = \langle 0,0|0,0\rangle_\delta = 1$
- $F_k = \langle \alpha_k, 0|\alpha_k, \beta_k\rangle_\delta = \langle \alpha_k, 0|0,0\rangle_\delta = 0$ with similar cross products also equal to zero.

Here, we use the notation of the bra/ket without any subscript for the field h_s and with a subscript δ for the field $h_s+\delta$, which will be followed through out this paper. We have also used the orthogonality of the basis states in the above steps. It is to be noted that within the entire gapless region including SLI and SLII phase, there is at least one mode for which $F_k = 0$ and hence fidelity in the entire gapless region is zero. This is also shown in Fig. 2.



FIG. 2: Variation of fidelity as a function of the staggered field h_s for two different values of J_3. The SLII phase exists only for $J_3 = 4$ but this phase is not captured by the fidelity. Fidelity only detects the gapped to gapless phase transition of the model which occurs at $h_s = J_3/2$.

IV. LOSCHMIDT ECHO

The ground state LE, defined as the overlap of the initial wavefunction, which is the ground state of the initial Hamiltonian, evolving under two slightly different Hamiltonians, is given as

$$LE = |\langle \Psi(h_s, t)|\Psi(h_s+\delta, t)\rangle|^2 \tag{10}$$

where $|\Psi(h_s, t)\rangle = e^{iH(h_s)t}|\Psi_G\rangle$ and $|\Psi(h_s+\delta, t)\rangle = e^{iH(h_s+\delta)t}|\Psi_G\rangle$, $|\Psi_G\rangle$ being the ground state for the Hamiltonian $H = H(h_s)$. In the momentum representation, as in the case of fidelity, one has to write an expression for the Loschmidt echo by taking the correct ground state for each mode. Thus,

$$\begin{aligned} LE &= \prod_k |\langle \Phi_k(h_s)|e^{iH(h_s+\delta)t}|\Phi_k(h_s)\rangle|^2 \\ &= \prod_k LE_k \end{aligned} \tag{11}$$

Let us calculate LE$_k$ for different possible ground states.

- If $|\Phi_k(h_s)\rangle = |\alpha_k, 0\rangle$

  $$LE_k = \left|\langle \alpha_k, 0|e^{iH(h_s+\delta)t}|\alpha_k, 0\rangle\right|^2 \tag{12}$$

  Since $|\alpha_k, 0\rangle$ is the eigenstate of $H(h_s)$ and not $H(h_s+\delta)$, we need to rewrite it in terms of $|\alpha_k, 0\rangle_\delta$. Using Eq. 6, we find that

  $$|\alpha_k, 0\rangle = \cos\eta_k|\alpha_k, 0\rangle_\delta - \sin\eta_k|0, \beta_k\rangle_\delta, \tag{13}$$

  where $2\eta_k = \theta_k(h_s) - \theta_k(h_s+\delta)$. Substituting the above transformation in Eq. 12, we get

  $$LE_k = 1 - \sin^2(2\eta_k)\sin^2\left(\frac{\Delta E(k)t}{2}\right) \tag{14}$$

  where $\Delta E(k) = E_+(h_s+\delta, k) - E_-(h_s+\delta, k) = 2\sqrt{(h_s+\delta)^2 + \cos^2(k/2)}$. Clearly, the three spin term has nothing to do with in ΔE and does not influence LE$_k$ though it will affect the magnitude of the dip in LE as shown in Fig. 3. This is because the number of $k-$ modes which influence the Loschmidt echo changes as J_3 changes.

- If $|\Phi_k(h_s)\rangle = |0,0\rangle$ or $|\alpha_k, \beta_k\rangle$, then $LE_k = 1$ as these are also the eigen vectors of the Hamiltonian $H_k(h_s+\delta)$.

From the above two points, it is clear that the LE will show a dip when ΔE is minimum at which $\sin^2 2\eta_k$ is nonzero. On the other hand, the ground state at $h_s+\delta = 0$ for the mode $k = \pi$ at which $\Delta E = 0$ is $|\alpha_k, \beta_k\rangle$ for $J_3 \neq 0$ and hence LE$_{k=\pi}$ is unity making this critical mode irrelevant in the dynamics. It is the other modes for which $E_+(k) > 0$ and $E_-(k) < 0$ which actually influence the behavior of the LE and is one of the interesting observations of the paper. Since the critical mode is not involved in the dynamics, various power-law scalings of LE are not present in this model. There is another way of looking at this point: Consider Eq. 5. As mentioned



FIG. 3: Variation of the LE with the staggered field h_s for different values of three spin interaction term.

before, $|0,0\rangle$ and $|a_k, b_k\rangle$ do not mix and we can concentrate on $|a_k, 0\rangle$ and $|0, b_k\rangle$ basis. In these two basis, H_k can be written as

$$\frac{J_3}{2}\cos k\hat{I} + h_s\sigma^z - \cos(k/2)\sigma^x \tag{15}$$

where second and third term combined together resembles the Landau -Zener Hamiltonian, though without the linear time dependence[25]. Clearly, the dynamics is governed by the minimum energy gap corresponding to the Landau Zener Hamiltonian[26] which occurs at $h_s = 0$ and hence only this point is detected by the LE. We call $h_s = 0$ and $k_c^d = \pi$ as the dynamical critical point and dynamical critical mode, respectively. Such an observation was also reported in the context of defect generation for a similar model where the authors claimed that the identity term only changes the phase of the evolving wave function and hence does not contribute to the defect generation[26].

We also studied the dependence of LE on system size and the coupling strength δ at the dynamical critical point $h_s + \delta = 0$. The LE decays exponentially with the system size N whereas it decays exponentially with δ^2 for both the regions, $|h_s| > J_3/2$ and $|h_s| < J_3/2$ as shown in Fig. 4. To understand this behavior, let us concentrate on S =log LE which is given by

$$S = \sum_k \log\left(1 - \sin^2 2\eta_k \sin^2 \frac{\Delta E_k t}{2}\right). \tag{16}$$



FIG. 4: Variation of log LE with system size N and the coupling strength δ for $h_s + \delta = 0$.

As mentioned before, ΔE_k never vanishes as the critical mode does not contribute to the decay of the LE. Since $\sin^2 2\eta_k$ is very small as it is the difference between two approximately equal angles (critical mode is not involved in the calculation), we can write

$$\begin{aligned} S &\simeq -\sum_k 4\eta_k^2 \sin^2(\frac{\Delta E_k t}{2}) \\ &\simeq -\sum_k \frac{\delta^2 \sin^2(t\cos k/2)}{\cos^2(k/2)} \end{aligned} \tag{17}$$

which explains the δ^2 dependence of S. It is difficult to understand the system size dependence of the LE since the critical mode does not contribute to the decay and hence no further simplifications can be done. But we do know that the mode closest to the dynamical critical mode $k_c^d = \pi$ for which $E_-(k) < 0$ and $E_+(k) > 0$ contribute maximum. Expanding Eq. 17 around $k = \pi$ for small time t, we find that

$$S = -\sum_k \delta^2 t^2 \propto N$$

and is shown in Fig. 4.

Though we considered a spin-1/2 model, such type of excitation spectrum, i.e., identity matrix and a Landau Zener type Hamiltonian, also exists in hard core Bose-Hubbard model in the presence of a period two superlattice[27]. We expect similar results to hold even in this hard core boson model which has various phases like superfluid, Mott insulator, hole vacuum and particle vacuum phase.

At the end, we note that there are a couple of works studying the fidelity and LE of models containing multispin interaction in Ising type Hamiltonians[28,29]. In Ref. 28, the effect of some complicated three spin interaction on LE is studied where the authors reported that a particular term does not effect the position of the dip in LE and changes only the sharpness of the decay. On the other hand, we have studied a completely different model with special types of phase transition points and presented arguments on why LE is not able to detect the gapped to gapless phase transition points. Also, the physics of our model is very different from the model studied in Ref. 28 due to the presence of staggered field. As a result of this, our model contains two types of quasiparticles making the handling of the problem very different from the other models studied till now in the context of fidelity or Loschmidt echo.

V. CONCLUSIONS

Usually, there is a sharp dip in the ground state fidelity or equivalently in the LE close to a quantum critical point. We have shown here an example of a unique quantum phase transition where neither fidelity nor LE can capture the entire phase diagram. While fidelity shows a dip at the boundary between the gapped to gapless phase transition and is unable to detect the two Fermi point to four Fermi point phase transition within the gapless region, the Loschmidt echo shows a completely different picture. It is able to detect only one special point in the entire phase diagram which is the point corresponding to the minimum of the Landau Zener type Hamiltonian and can not capture the critical points of the total Hamiltonian. This is because the dynamics is entirely governed by the Landau-Zener type Hamiltonian and the identity matrix containing the three spin term only adds a phase to the wave function evolution.

Acknowledgements: UD acknowledges fruitful discussions with Shraddha Sharma and thanks Amit Dutta for critically reading the manuscript.

---

[1] P. Zanardi and N. Paunkovic, Phys. Rev. E **74**, 031123 (2006).

[2] H-Q. Zhou and J. P. Barjaktarevic, J. Phys. A: Math. Theor. **41**, 412001 (2008).

[3] Shi-Jian Gu, Int. J. Mod. Phys. B **24**, 4371 (2010).

[4] S. Sachdev, *Quantum Phase Transitions*(Cambridge University Press, Cambridge, England,1999).

[5] B. K. Chakrabarti, A. Dutta and P. Sen, *Quantum Ising Phases and transitions in transverse Ising Models*, m41 (Springer, Heidelberg,1996).

[6] S. Yang, S. J. Gu, C. P. Sun and H-Q. Lin, Phys. Rev. A **78**, 012304 (2008); J. H. Zhao and H. Q. Zhou, arXiv:0803.0814

[7] P. Bounsante and A. Vezzani, Phys. Rev. Lett. **98**, 110601 (2007).

[8] H. T. Quan, Z. Song, X. F. Liu, P. Zanardi and C. P Sun, Phys. Rev. Lett. **96**, 140604 (2006).

[9] F. M. Cucchietti, S. F. Vidal and J. P. Paz, Phys. Rev. A **75**, 032337 (2007); D. Rossini, *et al*, Phys. Rev. A **75**, 032333 (2007); J. Zhang, *et al*, Phys. Rev. A **79**, 012305 (2009).

[10] V. Mukherjee, S. Sharma and A. Dutta, Phys. Rev. B **86**, 020301 (2012).

[11] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic Publishers, Dordrecht); A. Peres, Phys. Rev. A **30**, 1610 (1984).

[12] R. A. Jalabert, and H. M. Pastawski, Phys. Rev. Lett. **86**, 246 (2001)

[13] Z. P. Karkuszewski, C. Jarzynski, and W. H. Zurek, Phys. Rev. Lett. **89**, 170405 (2002)

[14] N. R. Cerruti, S. Tomsovic, Phys. Rev. Lett. **88**, 054103 (2002).

[15] F. M. Cucchietti, , D.A. R. Dalvit, J.P. Paz, and W.H. Zurek, Phys. Rev. A **95**, 105701 (2003).

[16] Q. Zheng, W. G. Wang, X. P. Zhang and Z. Z. Ren, Phys. Lett. A **372**, 5139 (2008); Q. Zheng, W. G. Wang, P. Q. Qin, P. Wang, X. P. Zhang and Z. Z. Ren, Phys. Rev. E **80**, 016214 (2009).

[17] J. F. Huan, Y. Li, J. Q. Liao. L. M. Kuang and C. P. Sun, Phys. Rev. A **80**, 063829 (2009).

[18] F. B. J. Buchkremer, R. Dumke, H. Levsen, G. Birkl and W. Ertmer, Phys. Rev. Lett. **85**, 3121 (2000).

[19] J. Zhang *et.al*, Phys. Rev. A **80**, 012305 (2009).

[20] C. M. Sánchez, P. R. Levstein, R. H. Acosta and A. K. Chattah, Phys. Rev. A **80**, 012328 (2009).

[21] M. Roger, J. H. Hetherington and J. M. Delrieu, Rev. Mod. Phys. **55**, 1 (1983).

[22] I. Titvinidze and G. I. Japaridze, Eur. Phys. J. B **32**, 383 (2003); A. A. Zvyagin and G. A. Skorobagatḱo, Phys. Rev. E **73**, 024427 (2006).

[23] E. Lieb, T. Schultz and D. Mattis, Ann. Phys.(N.Y.) **16** 37004 (1961).

[24] J.E. Bunder and R. H. McKenzie, Phys. Rev. B., **60**, 344, (1999).

[25] C. Zener, Proc. Roy. Soc. London Ser A **137**, 696 (1932); L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Non-relativistic Theory*, 2nd ed. (Pergamon Press, Oxford, 1965).

[26] D. Chowdhury, U. Divakaran and A. Dutta, Phys. Rev. E. **81**, 012101 (2010).

[27] I Hen, M. Iskin and M. Rigol, Phys. Rev. B **81**, 064503 (2010).

[28] H. L. Lian, Physica B **406**, 4278 (2011).

[29] H. L. Lian, D. P. Tian, Phys. Lett. A **375**, 3604 (2011); X. Liu, M. Zhing, H. Xu and P. Tong, J. Stat. Mech: Thoery and Experiment, P01003 (2012).